

$(U(p, q), U(p - 1, q))$ is a generalized Gelfand pair

Gerrit van Dijk

Received: 9 January 2008 / Published online: 2 September 2008
© The Author(s) 2008. This article is published with open access at Springerlink.com

Abstract Denote by $G = U(p, q)$ the orthogonal group of the sesqui-linear quadratic form $[x, y] = x_1 \bar{y}_1 + \cdots x_p \bar{y}_p - x_{p+1} \bar{y}_{p+1} - \cdots - x_{p+q} \bar{y}_{p+q}$ on \mathbb{C}^{p+q} and let $H_1 = U(p - 1, q)$ be the stabilizer of the first unit vector e_1 . Let $H_0 = U(1)$ and set $H = H_0 \times H_1$. Define the character χ_l of H by $\chi_l(h) = \chi_l(h_0 h_1) = h_0^l$ ($h_0 \in H_0, h_1 \in H_1$) where $l \in \mathbb{Z}$. Define the anti-involution σ on G by $\sigma(g) = \bar{g}^{-1}$. In this note we show that any distribution T on G satisfying $T(h_1 g h_2) = \chi_l(h_1 h_2) T(g)$ ($g \in G; h_1, h_2 \in H$) is invariant under the anti-involution σ . This result implies that (G, H_1) is a generalized Gelfand pair.

Keywords Generalized Gelfand pair · Unitary group · Anti-involution

Mathematics Subject Classification (2000) 22E30 · 43A85

1 Introduction

Let $G = U(p, q)$ be the orthogonal group of the sesqui-linear quadratic form on \mathbb{C}^n ($n = p + q$) given by

$$[x, y] = x_1 \bar{y}_1 + \cdots x_p \bar{y}_p - x_{p+1} \bar{y}_{p+1} - \cdots - x_{p+q} \bar{y}_{p+q},$$

and let $H_1 \simeq U(p - 1, q)$ be the stabilizer in G of the first unit vector e_1 of \mathbb{C}^n . We are going to show:

Theorem 1 *The pair (G, H_1) is a generalized Gelfand pair.*

Let us recall the definition of generalized Gelfand pair [6]. Let (π, \mathcal{H}) be any irreducible unitary representation of G on the Hilbert space \mathcal{H} . Then (G, H_1) is said to be a generalized Gelfand pair if $\dim \mathcal{H}_{-\infty}^{H_1} \leq 1$. Here $\mathcal{H}_{-\infty}^{H_1}$ is the space of H_1 -fixed vectors in $\mathcal{H}_{-\infty}$,

G. van Dijk (✉)
Mathematical Institute, Niels Bohrweg 1, 2333 CA Leiden, The Netherlands
e-mail: dijk@math.leidenuniv.nl

the (continuous) dual of the space \mathcal{H}_∞ of C^∞ -vectors. Equivalently one has: any unitary representation of G which can be realized on the space $D'(G/H_1)$ of distributions on G/H_1 decomposes multiplicity free.

In [1] another definition of Gelfand pair is used: for any irreducible admissible smooth Fréchet representation (π, E) of G one has E^{H_1} is at most one-dimensional. It has been confirmed by the authors that our results imply that the pair (G, H_1) is even a Gelfand pair in the sense of [1].

Theorem 1 is certainly true for $p = 1$, the Riemannian case. However in this case a much stronger result holds: any irreducible unitary representation of G decomposes, when restricted to the compact subgroup H_1 , multiplicity free. This was proved by Koornwinder in [4] using a criterion which reads as follows: any continuous function F on G satisfying $F(h_1 g h_1^{-1}) = F(g)$ for all $g \in G$, $h_1 \in H_1 \simeq \mathrm{U}(q)$, is invariant under the anti-involution σ of G defined by $\sigma(g) = \bar{g}^{-1}$ ($g \in G$).

2 A criterion

We shall formulate here a criterion (Theorem 2) which implies Theorem 1.

Let H be the subgroup $H = H_0 \times H_1$ of G where $H_0 = \mathrm{U}(1)$. So any element of H has the form

$$h = \begin{pmatrix} h_0 & 0 \\ 0 & h_1 \end{pmatrix}$$

with $h_0 \in H_0$, $h_1 \in H_1$. Define the character χ_l of H by $\chi_l(h) = h_0^l$, $l \in \mathbb{Z}$.

Let us denote by $d(\theta)$ the diagonal matrix in G with entries $e^{i\theta}$ ($\theta \in \mathbb{R}$). Clearly $d(\theta)$ belongs to the center of G ,

If (π, \mathcal{H}) is an irreducible unitary representation then by Schur's Lemma $\pi(d(\theta))$ acts as a scalar, say $e^{il\theta}$. Therefore the vectors in $\mathcal{H}_{-\infty}^{H_1}$ transform in the same way under $\pi_{-\infty}(d(\theta))$. This leads to the following criterion.

Theorem 2 *Let $l \in \mathbb{Z}$ be fixed. Any distribution T on G satisfying*

$$T(h_1 g h_2) = \chi_l(h_1 h_2) T(g) \quad (g \in G; h_1, h_2 \in H)$$

is invariant under the anti-involution of G defined by $\sigma(g) = \bar{g}^{-1}$.

This theorem implies Theorem 1, see [5].

3 Proof of Theorem 2

The idea of the proof was laid down in [2].

Let X_1 be the space defined by

$$X_1 = \{x \in \mathbb{C}^n : [x, x] = 1\}.$$

Then $X_1 \simeq G/H_1$, the isomorphism being given by $p_1 : g \mapsto g \cdot e_1$ ($g \in G$), where e_1 is the first standard unit vector in \mathbb{C}^n .

Let $D(X_1)$ denote the space of complex-valued C^∞ -functions on X_1 with compact support. The left G -action on X_1 induces a representation U of G on $D(X_1)$ and by inverse

transposition a representation U of G on $D'(X_1)$. We define

$$D'(X_1, l) = \{T \in D'(X_1) : U_h T = \chi_l(h) T \ (h \in H)\}.$$

This space is naturally isomorphic to the space considered in Theorem 2, the space of distributions T on G that satisfy the (formal) transformation rule

$$T(h_1 g h_2) = \chi_l(h_1 h_2) T(g) \quad (g \in G; h_1, h_2 \in H).$$

Let us introduce a map ξ which describes the H_1 -orbits on X_1 . Let x_1 be the first coordinate of $x \in X_1$. Consider the map $\xi : X_1 \rightarrow \mathbb{C}$ given by $\xi(x) = x_1$. It has the following properties:

- ξ is H_1 -invariant,
- ξ is real analytic,
- $\xi(x) = t$ ($t \in \mathbb{C}$) is an H_1 -orbit on X_1 if $t\bar{t} \neq 1$,
- ξ has no critical values: $\text{rank } d(\xi)(x) = 2$ if $t\bar{t} \neq 1$, $\text{rank } d\xi(x) = 1$ if $t\bar{t} = 1$.

Moreover we define $Q : X_1 \rightarrow [0, \infty)$ by

$$Q(x) = |\xi|^2.$$

We define also the following open subsets of X_1 :

$$X_1^0 = \{x \in X_1 : Q(x) < 1\}$$

$$X_1^1 = \{x \in X_1 : Q(x) > 0\}.$$

The map Q is left H -invariant, hence both sets are. Therefore we may define for $j = 0, 1$

$$D'(X_1^j, l) = \{T \in D'(X_1^j) : U_h T = \chi_l(h) T \ (h \in H)\}.$$

Since $X_1^0 \cup X_1^1 = X_1$, the map $T \mapsto (T|_{X_1^0}, T|_{X_1^1})$ defines a linear bijection from $D'(X_1, l)$ onto the set of pairs $(T_0, T_1) \in D'(X_1^0, l) \times D'(X_1^1, l)$ satisfying the matching condition

$$T_0|_{X_1^0 \cap X_1^1} = T_1|_{X_1^0 \cap X_1^1}.$$

We shall study such pairs of distributions. It is sufficient to study the subspaces $D'(X_1^j, l)$ separately, since X_1^j ($j = 0, 1$) are clearly σ -invariant.

The spaces are treated with different methods. We first deal with the space $D'(X_1^1, l)$ and use the now classical method of Faraut [3] with applying Tengstrand's results. For this purpose we introduce the space $X = X_1 / \sim$ where points x and y are identified if $x = \lambda y$ with $\lambda \in \mathbb{C}$, $|\lambda| = 1$. Let $x \mapsto \tilde{x}$ be the natural projection of X_1 onto X . Observe that $X \simeq G/H$. Set $p(g) = \widetilde{p_1(g)}$ ($g \in G$). Clearly Q is well-defined on X . In addition the point \tilde{e}_1 is an isolated non-degenerate critical point of Q on X . Set $X^1 = \widetilde{X_1^1}$. Notice that the map ξ does not vanish on X_1^1 . Therefore one readily sees that multiplication by ξ^l induces a bijection $D'(X^1)^H \rightarrow D'(X_1^1, l)$. Here $D'(X^1)^H$ denotes the space of H -invariant distribution on X^1 . We have the following results. There is a map $M : f \mapsto M_f$ which is surjective from $D(X^1)$ onto a space \mathcal{H}_η , defined by

$$M_f(t) = \int_X f(x) \delta(Q(x) - t) dx$$

where dx is a G -invariant measure on X . One calls $M_f(t)$ the average of f over the surface $Q(x) = t$. The space \mathcal{H}_η consists of functions φ on $(0, \infty)$ of the form

$$\varphi(t) = \varphi_0(t) + \eta(t)\varphi_1(t)$$

with $\varphi_0, \varphi_1 \in D((0, \infty))$ and η the “singularity function”

$$\eta(t) = Y(1-t)(1-t)^{n-2}$$

with Y the Heaviside function: $Y(t) = 1$ if $t \geq 0$, $Y(t) = 0$ if $t < 0$.

Moreover, the transpose M' of M is injective from \mathcal{H}'_η to $D'(X^1)$ with image $D'(X^1)^H$. From this we conclude that any bi- H -invariant distribution T on $p^{-1}(X^1) \subset G$ satisfies $T = T^\sigma$. The proof is standard, see [5]. For convenience we repeat the argument here. Fix Haar measures dg on G and dh on H in such a way that $dg = dx dh$. For $f \in D(G)$ set

$$f^0(x) = \int_H f(gh) dh \quad (x = p(g) \in X).$$

Given a bi- H -invariant distribution T on $p^{-1}(X^1) \subset G$ there is a unique H -invariant distribution T_1 on X^1 satisfying $\langle T, f \rangle = \langle T_1, f^0 \rangle$ ($f \in D(p^{-1}(X^1))$), and conversely. Extend the function Q from X to G by $Q(g) = \|g \cdot e_1, e_1\|^2$. To show that T is σ -invariant, it is sufficient to show that

$$M_{[(f^\sigma)^0]} = M_{f^0}$$

for all $f \in D(p^{-1}(X^1))$. This is easily checked. For all continuous functions F on $(0, \infty)$ one has

$$\begin{aligned} \int_0^\infty F(t) M_{[(f^\sigma)^0]}(t) dt &= \int_{X^1} F(Q(x)) (f^\sigma)^0(x) dx \\ &= \int_G F(Q(g)) f^\sigma(g) dg = \int_G F(Q(\sigma(g))) f(g) dg. \end{aligned}$$

Since $Q(g) = Q(\sigma(g))$ ($g \in G$) we get the result.

Since the function ξ , extended to G by

$$\xi(g) = [g \cdot e_1, e_1],$$

is also σ -invariant, we get that any distribution T on $p_1^{-1}(X_1^1)$ satisfying

$$T(h_1 g h_2) = \chi_l(h_1 h_2) T(g) \quad (g \in G; h_1, h_2 \in H)$$

is σ -invariant.

We now consider the other space $D'(X_1^0, l)$. Here we use the map ξ . Recall that ξ is a submersion from X_1^0 onto $U = \{z \in \mathbb{C}; |z| < 1\}$. Its level sets are H_1 -orbits on X_1^0 . So we have the following lemma.

Lemma 1 *The natural pull-back map*

$$\xi^* : D'(U) \rightarrow D'(X_1^0)$$

is injective with image $D'(X_1^0)^{H_1}$.

It is now easy to show, as before, that any distribution T on $p^{-1}(X_1^0) \subset G$ satisfying

$$T(h_1 g h_2) = \chi_l(h_1 h_2) T(g) \quad (g \in G; h_1, h_2 \in H)$$

is again σ -invariant, since ξ (extended to G) is.

This completes the proof of Theorem 2.

Open Access This article is distributed under the terms of the Creative Commons Attribution Noncommercial License which permits any noncommercial use, distribution, and reproduction in any medium, provided the original author(s) and source are credited.

References

1. Aizenbud, A.A., Gourevitch, D., Sayag, E.: $(O(V \otimes F), O(V))$ is a Gelfand pair for a quadratic space V over a local field F , arXiv: 0711.1471v1 [math.RT] (2007)
2. Charchov, I.: Harmonic Analysis On Line Bundles Over Complex Hyperbolic Spaces. PhD Thesis, Leiden University (2000)
3. Faraut, J.: Distributions sphériques sur les espaces hyperboliques. J. Math. Pures Appl. **58**, 369–444 (1979)
4. Koornwinder, T.H.: A note on the multiplicity free reduction of certain orthogonal and unitary groups. Indag. Math. **44**, 215–218 (1982)
5. van Dijk, G.: On a class of generalized Gelfand pairs. Math. Z. **193**, 581–593 (1986)
6. van Dijk, G., Poel, M.: The Plancherel formula for the pseudo-Riemannian space $SL(n, \mathbb{R})/GL(n - 1, \mathbb{R})$. Compos. Math. **58**, 371–397 (1986)